

Probing nonperturbative QED with optimally focused laser pulses

A. Gonoskov,^{1,2,*} I. Gonoskov,^{1,†} C. Harvey,^{3,‡} A. Ilderton,^{1,4,§}
A. Kim,^{2,¶} M. Marklund,^{1,4,**} G. Mourou,^{5,6,††} and A. M. Sergeev^{2,‡‡}

¹*Department of Physics, Umeå University, SE-90187 Umeå, Sweden*

²*Institute of Applied Physics, Russian Academy of Sciences, Nizhny Novgorod 603950, Russia*

³*Centre for Plasma Physics, Queen's University Belfast, BT7 1NN, UK*

⁴*Department of Applied Physics, Chalmers University of Technology, SE-41296 Gothenburg, Sweden*

⁵*Institut de la Lumière Extrême, ENSTA, Palaiseau, France*

⁶*University of Nizhny Novgorod, Nizhny Novgorod 603950, Russia*

We study nonperturbative pair production in intense, focused laser fields called ‘*e*-dipole’ pulses. We address the conditions required, such as the quality of the vacuum, for reaching high intensities without initiating beam-depleting cascades, the number of pairs which can be created, and experimental detection of the created pairs. We find that *e*-dipole pulses offer an optimal method of investigating nonperturbative QED.

Understanding quantum field theory in the nonperturbative regime remains a challenging theoretical and experimental issue. Recent advances in technology have spurred interest in the possibility of using intense lasers to probe quantum vacuum phenomena, including nonperturbative electron-positron pair production [1, 2]. Such studies are complementary to those of, e.g., heavy ion collisions, with the possible advantage that laser systems provide a more controlled environment.

It is well known that nonperturbative pair creation is strongly suppressed below the Sauter-Schwinger limit $E_S = m^2 c^3 / e \hbar \simeq 10^{18}$ V/m [3, 4], and while field-strengths of this scale are typical of QED, creating them on a macroscopic (laboratory) scale will remain out of our reach for the foreseeable future. Various mechanisms for stimulating pair production have therefore been proposed; these include colliding an intense laser pulse with high energy photons [5], electrons [6], or other laser pulses [7–9]. The number of pairs which can be created in the collision of laser pulses is sensitive not just to the field amplitude [10], but also to the field structure [11]. The same applies to particle dynamics in general. In simulating intense laser-matter interactions it is therefore necessary to employ realistic field models.

It was pointed out in [12], see also [13], that there is a potential obstacle to reaching the Sauter-Schwinger limit, or high intensities in general. In an experiment, the presence of stray particles resulting from imperfect vacuum, or the generation of hard radiation, could trigger an avalanche of particle production. This perturbatively triggered cascade could hinder any observation of nonperturbative effects, both via beam depletion and by producing a background which swamps the signals of interest. Hence, it is important to understand the conditions leading to cascades, and what backgrounds they produce.

In this paper we consider Schwinger pair production in ‘*e*-dipole’ pulses [14]. These are exact, closed form solutions of Maxwell’s equations in vacuum. They exhibit optimal focusing efficiency, i.e. they produce the highest

possible peak field-strength for a given input power [15] and describe genuinely pulsed fields, having finite energy and finite extent in all four spacetime directions. We will show in a separate article that it is possible, using a new focusing system, to create a close approximation to an *e*-dipole pulse, hence pushing laser systems to near-optimal conditions [16]. Here we focus on exact *e*-dipole pulses in order to provide analytic estimates; we will show that the structure of *e*-dipole pulses may reduce the chances of cascade formation, thus allowing laser experiments to approach higher intensity regimes.

This paper is organised as follows. We begin by introducing *e*-dipole pulses. We then analyse particle motion in such pulses, showing that stray particles in an experiment are not driven into the high intensity part of the field, provided one begins with a vacuum of sufficient quality. We estimate the level of vacuum required to both keep particles away from the focus, and to limit the generation of hard photons into the focus. We then calculate the number of pairs which could be produced by our pulse and compare with other pulse models in the literature. The behaviour of electron-positron pairs post-creation is then analysed before we conclude.

Dipole pulses:– There is a limit to the focusing efficiency of a given laser system. Consider first an unfocused, broad laser pulse which can be regarded as nearly monochromatic. The peak field strength E_0 which can be obtained by focusing monochromatic light of wavelength λ and cycle-averaged power P is bounded above; defining $\mathbb{P} = P / (4\pi\epsilon_0 c)$, the peak field obeys [15]

$$E_0 \leq \frac{8\pi}{\sqrt{3}} \frac{\sqrt{\mathbb{P}}}{\lambda} \simeq 14.51 \frac{\sqrt{\mathbb{P}}}{\lambda}. \quad (1)$$

The focused fields which saturate this upper bound are called *e*-dipole pulses, see [14] for full details. *E*-dipole pulses take their name from structural similarities with dipole fields, but do not contain singularities. They describe a converging pulse of light, with the ideal case of 4π focusing. (The formation of this focused pulse can be

pictured as the reverse process of emission from a dipole antenna.) The e -dipole field has the following form. Let $R^2 = x^2 + y^2 + z^2$ and define the vector $\mathbf{Z}$ by

$$\mathbf{Z} = \hat{\mathbf{z}} \frac{d}{R} [g(t + R/c) - g(t - R/c)] , \quad (2)$$

in which the ‘driving function’ g is arbitrary and the ‘virtual dipole moment’ d is a constant. (Using the fact that e -dipole pulse can be generated by focusing laser fields [14], we will shortly relate g and d to input laser parameters.) The electromagnetic fields of an e -dipole pulse are then given in terms of $\mathbf{Z}$ by

$$\mathbf{E} = -\nabla \times \nabla \times \mathbf{Z}, \quad \mathbf{B} = -\frac{1}{c^2} \nabla \times \dot{\mathbf{Z}} . \quad (3)$$

These fields are exact solutions to Maxwell’s equations in vacuum. It is easily confirmed that there is no singularity at $R = 0$, the focus point. Far from the focus, the electric and magnetic field amplitudes of the e -dipole pulse are proportional to $dR^{-1}\ddot{g}(t \pm \frac{R}{c})$, and have an angular distribution proportional to $\sin^2 \theta$, with θ the polar angle. In the focus one has $\mathbf{E}(0, t) = \hat{\mathbf{z}} \frac{4d}{3c^3} \ddot{g}(t)$, $\mathbf{B}(0, t) = 0$. In general we are interested in pulses with, say, Gaussian frequency spreads as for the following driving function

$$g(\tau) = e^{-(\tau^2/D^2) \ln 4} \sin(\omega\tau) , \quad (4)$$

in which ω is the central frequency and D is the full-width-half-maximum duration (i.e. the intensity $\sim E^2$ drops to 1/2 its peak value at $t = \pm D/2$). We refer to such a pulse as ‘quasi-Gaussian’ since, far from the focus, the envelope is $\ddot{g}$, which has the same Gaussian frequency spread as g . It follows from (3) that such pulses are compactly supported in all four space-time dimensions, which is one of the advantages of e -dipole pulses over other field models in the literature; they describe genuine pulses, without sacrificing Maxwell’s equations.

The virtual dipole moment d can be expressed as a function of input energy, or power, by using energy conservation in the far-field. To illustrate, consider a monochromatic driving field, wavelength λ and cycle-averaged power $P := \frac{1}{2} P_{peak}$. The peak focused field strength E_0 , incoming power, and d are then related by

$$E_0 = \frac{4}{3} \frac{\omega^3}{c^3} d = \frac{8\pi}{\sqrt{3}} \frac{\sqrt{P}}{\lambda} , \quad (5)$$

Note that (5) is optimal, saturating the upper bound in (1). Similar relations can be evaluated for the quasi-Gaussian pulse [14]. This leads to a simple measure for comparing different pulse models. We define the focusing efficiency parameter f_E via $E_0 = f_E \frac{\sqrt{P}}{\lambda}$; this relates the incoming power P to peak field-strength E_0 . In order to compare with other models, this definition applies to fields which are monochromatic in the far field region. In Table I we compare the focusing parameters for

	Two NF beams [10]	Fedotov [17]	e -dipole
f_E	$8\pi\Delta \rightarrow 2.51$	$2\pi\sqrt{\frac{5}{3}} \simeq 8.11$	$\frac{8\pi}{\sqrt{3}} \simeq 14.51$

TABLE I. Focussing efficiency f_E (input power to peak electric field) for three beam configurations. We use $\Delta = 0.1$ to compare with the literature ($\Delta \ll 1$ is required [10]). The e -dipole pulse is most efficient.

three fields: the theoretical maximum from Fedotov’s solution [17], two colliding Narozhny-Fofanov beams [18] and the e -dipole field (with monochromatic g). The e -dipole pulse is most efficient.

Given a particular pulse model, one can increase (decrease) the focal electric (magnetic) field-strength by colliding several such pulses [9]. This increases the total focusing efficiency and is why, in Table I, we compare with two colliding beams from [18], as advocated for pair production in [19]. Using two pulses eliminates the focal magnetic field of the single pulse solution, and increases the peak field strength by a factor of $\sqrt{2}$. The potential increase using this method is limited, though; the behaviour of the efficiency depends strongly on the focusing geometry, due to the requirement that the total angle of incoming radiation cannot exceed 4π . As multiple beams start to overlap, they interfere, and the peak field strength is reduced. In particular, the ‘ $\sqrt{n}$ increase in field strength from n pulses’ is limited, for fixed and reasonable focusing angles, to rather small n .

Interaction with particles within the chamber:— A potential barrier to the observation of nonperturbative pair production is, simply, perturbative pair production, which is well understood. This and other processes can be initiated by particles within the imperfect vacuum of a chamber prepared for a laser experiment. Due to the effect of the ponderomotive force, an incoming laser pulse can ‘grab’ stray particles and accelerate them into the focus, leading to the emission of high energy photons which create pairs, which emit photons which create further pairs. The products of this cascade can obscure any nonperturbatively produced pairs, or the resulting beam depletion can be so great as to prevent sufficiently high intensities for Schwinger pair creation being reached [12]. If, though, we could guarantee that within the focal volume there were no high energy electrons and photons initiating perturbative processes (or rather, that such events had a low probability), then we could focus beams to the intensities required to trigger nonperturbative pair creation, without first initiating cascades, which is the experimental result of interest.

To investigate this prospect, one can determine the volume of space from which electrons can initiate cascades, as a function of the initial electron density (initial level of vacuum), and then require that this volume contain less than one particle. We therefore carried out a simulation of a large number of initially uniformly distributed




FIG. 1. Simulation results for electron motion under the influence of an e -dipole pulse. *Left scale*: initial number (N_0) and final average number (N) of electrons dragged into a sphere of radius R about the focus, as a function of R . Initial particle density 10^5 cm^{-3} . *Right scale*: maximum values of χ and ξ , taken over all electrons outside the sphere with radius R .

electrons moving (with radiation reaction accounted for by the Landau-Lifshitz equation [20]) in the fields of a converging e -dipole pulse, wavelength 810 nm, pulse duration 30 fs (consistent with candidate Ti-sapphire technology for future laser facilities [21, 22]) and total power $P = 1000 \text{ PW}$ (as is estimated, below, to be required for pair creation, see also Table II). Laser experiments are now routinely performed in technical vacua of stray molecule density 10^{-8} – 10^{-9} mbar, and lower pressures are accessible. We will assume an initial density of 10^5 cm^{-3} , equivalent to a pressure of 10^{-12} mbar. This figure is at the limit of what is achievable today, so it is a reasonable assumption for future facilities. The incoming laser pulse will ionise stray molecules and produce electrons. Now, we are interested only in whether electrons are pushed into the focus or not, based on their initial position. Thus, for the determination of the required vacuum quality, we can associate every molecule with a single electron, independent of the level of ionisation. We therefore take the initial electron density to be 10^5 cm^{-3} .

The results of our simulation are shown in Fig. 1. We show here the average number of electrons initially distributed within (N_0 , calculated from the initial density), or dragged into (N), a sphere of radius R around the focus, as a function of R . We see that for the chosen initial density, the final number N of particles drops below one at a radius of around $R^* = 30 \mu\text{m}$; this is much larger than the focus of the dipole pulse.

We immediately perform two checks on this result. First, we should only base our conclusions on data from regions of space in which quantum effects are negligible. Second, though particles remain *outside* the focus, they may still be accelerated such that they emit hard pho-

tons (capable of initiating cascades). In order to verify our result, we tracked in our simulation two parameters, χ and ξ . The first, χ , is the quantum efficiency parameter of the particles, calculated along their orbits [23]. This parameter estimates the importance of quantum effects and is, for a particle of momentum p_μ in a field with energy-momentum tensor $T_{\mu\nu}$ [24]

$$\chi = \frac{e\hbar}{m^3 c^4} \sqrt{p_\mu T^{\mu\nu} p_\nu} \approx \gamma \frac{E}{E_s}. \quad (6)$$

In the electron's instantaneous rest frame, χ is the ratio of the energy seen by the electron, to the electron's rest energy. As χ approaches unity, quantum effects start to become important; in high intensity fields χ is the input parameter for determining the probability of QED processes [23]. The second parameter, ξ , estimates the energy of emitted photons, in ratio to twice the rest mass of an electron. Recall that the typical energy $\hbar\omega_s$ of photons emitted by an ultra-relativistic electron is given in the synchrotron approximation by $\omega_s = \gamma^2 (eE/mc)$ [20]. We therefore define

$$\xi = \frac{\hbar\omega_s}{2mc^2} \approx \frac{1}{2} \gamma \chi, \quad (7)$$

So, for $\xi \ll 1$ we should be able to neglect the emission of hard photons with energy approaching $2mc^2$.

Returning to our results, Fig. 1 also shows the *maximum* value (over all electrons) of χ and ξ *outside* the sphere of radius R . We can conclude first that the classical description is valid ($\chi \ll 1$) at least outside the sphere of radius R^* . Hence, our estimate of the density increase within this sphere (due to dragging from outside it) can be trusted. Second, since $\xi < 1$, particles outside the sphere do not generate photons suitable for initiating pair creation within the sphere. Hence we can conclude from Fig. 1 that an initial density of less than 10^5 cm^{-3} leads to an average of well less than one particle existing in or entering into the sphere of radius R^* , and an absence of photons sufficiently hard to create pairs within the sphere.

Our estimate for the required initial density is overly cautious: more accurate estimates which include data on, e.g. the emission direction of radiation (our simulations show for example that hard photons are typically emitted away from, rather than toward, the focus [16]) could significantly reduce the stated requirements.

Pair production:– The quantities relevant for pair production are not the electric or magnetic field strengths themselves, but rather the local Lorentz invariants [4]

$$\mathcal{S} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu} \quad \text{and} \quad \mathcal{P} = -\frac{1}{4} \tilde{F}_{\mu\nu} F^{\mu\nu}. \quad (8)$$

In an e -dipole pulse we have $\mathcal{P} = \mathbf{E} \cdot \mathbf{B} = 0$, i.e. the electric and magnetic fields are orthogonal in all space. Near the focus of the pulse, the invariant $\mathcal{S}$ is positive

	$\lambda = 1 \mu\text{m}$	$\lambda = 0.8 \mu\text{m}$	$\lambda = 0.4 \mu\text{m}$
1660 PW	1 (5.5 kJ)	10^3 (4.4 kJ)	10^{10} (2 kJ)
1120 PW	10^{-4} (3.7 kJ)	1 (3 kJ)	10^8 (1.5 kJ)
320 PW	10^{-23} (1.1 kJ)	10^{-14} (0.85 kJ)	1 (0.43 kJ)

TABLE II. The number of pairs produced (order of magnitude only) as a function of wavelength and power. Bracketed figures give the incoming energy in one cycle. Parameters are chosen so that the diagonal values exhibit the threshold (average) power required to create a single pair in one cycle.

(the electric field ‘dominates’) and the pulse is capable of pair production [3, 4, 25]. In general, the key to producing a significant number of pairs is maximising the relevant invariant. This can be done by maximising the electric field and eliminating the magnetic field in the focus, which in practice is achieved (as above) by the use of counter-propagating pulses [9, 19]. Define a second focusing parameter, f_ϵ , via $\epsilon = f_\epsilon \frac{\sqrt{P}}{E_S \lambda}$. This relates incoming power to peak invariant ϵ . For two NF beams one finds $f_\epsilon = 8\pi\Delta \simeq 2.51$ for $\Delta = 0.1$ (a single NF beam has $f_\epsilon = 16\pi\Delta^2$ due to a nonzero focal magnetic field), while the e -dipole pulse has $f_\epsilon = f_E = \frac{8\pi}{\sqrt{3}} \simeq 14.5$. Again, the dipole pulse gives the most efficient ‘conversion’ of input power into invariant ϵ , and so we expect the e -dipole pulse to be capable of producing significantly more pairs than is possible in other pulses.

To calculate the number of produced pairs, we use the locally-constant-field estimate [10], based on [25]. For a field in which $\mathcal{P} \equiv 0$ we have

$$N_{\text{pairs}} = \frac{1}{4\pi^3 \lambda_c^4} \int d^4x \epsilon^2(x) \exp\left[-\frac{\pi}{\epsilon(x)}\right], \quad (9)$$

where $\epsilon = \sqrt{\mathcal{S} + |\mathcal{S}|}/E_S$, and $\lambda_c = \hbar/mc$ is the (reduced) Compton wavelength. Since the main contribution to N_{pairs} comes from the focal region (which in a quasi-Gaussian is localised both in space and time), and in order to compare with other pulse models (which are typically monochromatic in the far-field), we temporarily drop the envelope in (4), setting $D = \infty$, and quote only the contribution to N_{pairs} from a single cycle.

The number of pairs created is shown in Table II, as a function of the driving field’s wavelength and (average) input power. Our results demonstrate that pair production can be observed by using several kJ of total energy, at optical wavelength: for a driving pulse with $\lambda = 0.8\mu\text{m}$, the threshold power required for pair creation is around $P = 1120 \text{ PW}$, and the corresponding peak electric field is $E_0 \simeq 0.08E_S$. We see therefore that large numbers of pairs can be produced well below the Schwinger limit, compare [9, 17].

All predictions of the pair yield based on (9) should of course be interpreted as a measure of how easy it is to *produce* a pair, given an initial power and frequency, rather than a number of pairs. The reason for this is that

once a pair is produced, other processes can occur and either increase or decrease the number of pairs able to escape the pulse and be detected. In the former category is cascade formation [13, 26], in which the pairs are accelerated and emit hard photons which create further pairs, and so on [11, 12]. In the latter category are processes such as pair annihilation, though the annihilation cross section in intense fields is typically much smaller than that of cascade-generating processes [23, 27]. It is therefore important to address how one might best detect pair production events. We turn to this now, by analysing the behaviour of the created pairs.

Post-creation behaviour:— Fig. 2 shows typical trajectories, and χ values, for positrons born at rest near the focus of a $D = 30 \text{ fs}$ e -dipole pulse. Due to the intricate field structure, the trajectories depend sensitively on where the particles are created. The majority of particles, with a typical trajectory shown in red in Fig. 2, have large χ -values, implying that the depicted motion will receive significant quantum corrections. We can see though that if the particles leave the focus, their χ drops quickly to below one and classical predictions become accurate. Our simulations show that radiation reaction is responsible for recirculating particles back into the focus, and has an added effect of keeping the particles ‘in phase’ with the field intensity gradient, which is the cause of their large χ values. Recirculation returns the particles to the quantum regime, increasing the possibility of additional QED processes occurring, as predicted in [13]. Hence, even at intensities well below the Sauter-Schwinger limit, the dipole pulse becomes a testing ground not for classical electrodynamics, but for QED.

The behaviour of particles born very close to the focus is different. Those particles simply exit the pulse, without oscillation, moving almost parallel to the z -axis. Further, our simulations show that these particles are ‘out of phase’ with the field, so that while they have high γ factors, they also have $\chi < 1$, and so can be analysed classically. These particles will emit radiation in a small cone about the z -axis. We therefore suggest the detection of particles exiting in the z -direction as a possible candidate for direct measurement of the produced pairs.

Conclusions:— We have considered nonperturbative pair production in the fields of an e -dipole pulse. For sufficient levels of vacuum, our estimates (which may be considered cautious, given the inevitable improvements in vacuum technology which will be commensurate with laser development) show that stray particles are not driven into the most intense part of the field, thus reducing the chances of cascade initiating and beam depletion. This allows higher focal intensities to be reached, and allows us to consider the possibility of experimentally measuring nonperturbative pair creation, for which a total power of 1000 PW is estimated to be necessary. This corresponds to field-strengths well below the Schwinger




FIG. 2. Trajectories [left] and χ parameter [right] for particles created in a 30 fs, 1100 PW pulse. Initial positions: $x = y = z = 10^{-3}\lambda$ (red), $x = y = z = 10^{-4}\lambda$ (green), $x = y = z = 10^{-5}\lambda$ (blue).

limit.

We thank the Joint Supercomputer Centre of RAS and the Swedish National Infrastructure for Computing (SNIC) for provided supercomputer sources. A.I. thanks C. Murphy for useful discussions. The authors are supported by the Ministry of Education and Science of the Russian Federation, Agreement No. 11.G34.31.0011 (A.G., G.M., A.S.), the Swedish Research Council, contract 2011-4221 (A.I.), the European Research Council contract 204059-QPQV (A.I., M.M.) and EPSRC, grant EP/I029206/1-YOTTA (C.H.).

* arkady.gonoskov@physics.umu.se

† ivan.gonoskov@physics.umu.se

‡ christopher.harvey@qub.ac.uk

§ anton.ilderton@physics.umu.se

¶ kim@ufp.appl.sci-nnov.ru

** mattias.marklund@physics.umu.se

†† gerardmourou@gmail.com

‡‡ ams@ufp.appl.sci-nnov.ru

- [1] G. V. Dunne, Eur. Phys. J. D **55** (2009) 327
- [2] A. Di Piazza *et al.*, Rev. Mod. Phys. **84** (2012) 1177.
- [3] F. Sauter, Z. Phys. **69**, 742 (1931).
- [4] J. S. Schwinger, Phys. Rev. **82**, 664 (1951).
- [5] V. N. Baier and V. M. Katkov, Phys. Lett. A **374** (2010) 2201.
- [6] C. Bamber *et al.*, Phys. Rev. D **60**, 092004 (1999).
- [7] R. Schützhold, H. Gies and G. Dunne, Phys. Rev. Lett. **101**, 130404 (2008)
- [8] G. V. Dunne, H. Gies and R. Schutzhold, Phys. Rev. D **80**, 111301 (2009)
- [9] S. S. Bulanov, V. D. Mur, N. B. Narozhny, J. Nees and V. S. Popov, Phys. Rev. Lett. **104** (2010) 220404.
- [10] S. S. Bulanov, N. B. Narozhny, V. D. Mur and V. S. Popov, Phys. Lett. A **330** (2004) 1.
- [11] S. S. Bulanov, T. Z. Esirkepov, A. G. R. Thomas, J. K. Koga and S. V. Bulanov, Phys. Rev. Lett. **105** (2010) 220407.
- [12] A. M. Fedotov, N. B. Narozhny, G. Mourou and G. Korn, Phys. Rev. Lett. **105** (2010) 080402.
- [13] N. V. Elkina, A. M. Fedotov, I. Y. Kostyukov, M. V. Legkov, N. B. Narozhny, E. N. Nerush and H. Ruhl, Phys. Rev. ST Accel. Beams **14** (2011) 054401.
- [14] I. Gonoskov, A. Aiello, S. Heugel, G. Leuchs, Phys. Rev. A **86** (2012) 053836.
- [15] I. M. Bassett, Optica Acta: Int. J. Optics **33** (1986) 279.
- [16] In preparation.
- [17] A. M. Fedotov, Laser Physics **19** (2009) 214.
- [18] N. B. Narozhny, M. S. Fofanov, JETP **117** (2000) 867.
- [19] S. S. Bulanov, N. B. Narozhny, V. D. Mur and V. S. Popov, JETP **102** (2006) 9.
- [20] L. D. Landau, E. M. Lifshitz, *The classical theory of fields*, 1975, Elsevier, Oxford.
- [21] <http://www.extreme-light-infrastructure.eu/>.
- [22] <http://www.xcels.iapras.ru>.
- [23] V. I. Ritus, J. Russian Laser Research **6** (1985) 497.
- [24] T. Heinzl and A. Ilderton, Opt. Commun. **282** (2009) 1879.
- [25] W. Heisenberg, H. Euler, Z. Phys., **33**, 714, (1936).
- [26] A. R. Bell and J. G. Kirk, Phys. Rev. Lett. **101** (2008) 200403.
- [27] A. I. Voroshilo, E. A. Padusenko and S. P. Roshchupkin, Laser Phys. **20** (2010) 1679.